

H1N1 flu virus

Your questions answered

What is the novel H1N1 virus?

The 2009 H1N1 flu virus (initially known as swine flu) is a respiratory disease caused by type A influenza viruses that causes regular outbreaks in pigs. People do not normally get swine flu, but human infections can and do happen. Swine flu viruses have been reported to spread from person-to-person, but in the past, this transmission was limited and not sustained beyond three people.

The novel H1N1 flu has two genes from flu viruses that normally circulate in pigs in Europe and Asia and bird (avian) genes and human genes. Scientists call this a "quadruple reassortant" virus.

Are there human infections of H1N1?

Yes. Human infections with the new H1N1 virus are ongoing in the United States. Most people who have become ill with this new virus have recovered without requiring medical treatment.

CDC routinely works with states to collect, compile and analyze information about influenza, and has done the same for the new H1N1 virus since the beginning of the outbreak.

Is H1N1 contagious?

CDC has determined that novel H1N1 virus is highly contagious and is spreading from human to human.

What are the signs and symptoms of H1N1 flu in people?

The symptoms of novel H1N1 flu virus in people include fever, cough, sore throat, runny or stuffy nose, body aches, headache, chills and fatigue. A significant number of people who have been infected with this virus also have reported diarrhea and vomiting. Severe illnesses and death has occurred as a result of illness associated with this virus.

How severe is illness associated with H1N1?

Illness with the new H1N1 virus has ranged from mild to severe. While most people who have been sick have recovered without needing medical treatment, hospitalizations and deaths from infection with this virus have occurred.

The priority groups to receive the H1N1 vaccination initially are:

- All people 6 months through 24 years of age
- People who live with or care for children younger than 6 months of age
- All pregnant women
- Health care and emergency services personnel, and

What Can **YOU** Do?

- People ages 25 through 64 years who have health conditions associated with higher risk of medical complications from influenza.

Adults 64 and older do not appear to be at increased risk of novel H1N1-related complications thus far. About one-third of adults older than 60 may have antibodies against this virus. It is unknown how much, if any, protection may be afforded against novel H1N1 flu by any existing antibody.

How does H1N1 flu compare to seasonal flu in terms of its severity and infection rates?

With seasonal flu, we know that seasons vary in terms of timing, duration and severity. Seasonal influenza can cause mild to severe illness, and at times can lead to death. Each year, in the United States, on average 36,000 people die from flu-related complications and more than 200,000 people are hospitalized from flu-related causes. Of those hospitalized, 20,000 are children younger than 5 years old. More than 90 percent of deaths and about 60 percent of hospitalization occur in people older than 65.

When the novel H1N1 outbreak was first detected in mid-April 2009, CDC began working with states to collect, compile and analyze information regarding the novel H1N1 flu outbreak, including the numbers of confirmed and probable cases and the ages of these people.

The information analyzed by CDC supports the conclusion that novel H1N1 flu has caused greater disease burden in people younger than 25 years of age than older people. At this time, there are few cases and few deaths reported in people older than 64 years old, which is unusual when compared with seasonal flu.

However, pregnancy and other previously recognized high risk medical conditions from seasonal influenza appear to be associated with increased risk of complications from this novel H1N1. These underlying conditions include asthma, diabetes, suppressed immune systems, heart disease, kidney disease, neurocognitive and neuromuscular disorders and pregnancy.

How long can an infected person spread H1N1 to others?

People infected with seasonal and novel H1N1 flu shed virus and may be able to infect others from 1 day before getting sick to 5 to 7 days after. This can be longer in some people, especially children and people with weakened immune systems and in people infected with the new H1N1 virus.

How can I protect myself from getting sick and spreading the flu to others?

Right now, there is no vaccine for the H1N1 virus.

However, a novel H1N1 vaccine is currently in production and may be ready for the public in the fall. As always, a vaccine will be available to protect against seasonal influenza

There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza.

Take these everyday steps to protect your health:

- Cover your nose and mouth with a tissue when you cough or sneeze.

MORE

Throw the tissue in the trash after you use it.

- Wash your hands often with soap and water, especially after you cough or sneeze. Alcohol-based hand cleaners* are also effective.
- Avoid touching your eyes, nose or mouth. Germs spread this way.
- Try to avoid close contact with sick people.
- If you are sick with flu-like illness, CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from
- others as much as possible to keep from making others sick.

Other important actions that you can take are:

- Follow public health advice regarding school closures, avoiding crowds and other social distancing measures.
- Be prepared in case you get sick and need to stay home for a week or so; a supply of over-the-counter medicines, alcohol-based hand rubs,* tissues and other related items might could be useful and help avoid the need to make trips out in public while you are sick and contagious.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children, emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- Bluish or gray skin color
- Not drinking enough fluids
- Severe or persistent vomiting
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough.

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting
- Flu-like symptoms improve but then return with fever and worse cough.

If you are sick with flu-like illness, CDC recommends that you stay home for at least 24 hours after your fever is gone except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from others as much as possible. Cover your mouth and nose with a tissue when coughing or sneezing. Put your used tissue in the waste basket. Then, clean your hands, and do so every time you cough or sneeze.

Employees who are well but who have an ill family member at home with novel H1N1 flu can go to work as usual. These employees should monitor their health every day, and take everyday precautions including washing their hands often with soap and water, especially after

they cough or sneeze. Alcohol-based hand cleaners are also effective.* If they become ill, they should notify their supervisor and stay home. Employees who have an underlying medical condition or who are pregnant should call their health care provider for advice, because they might need to receive influenza antiviral drugs to prevent illness. For more information please see General Business and Workplace Guidance for the Prevention of Novel Influenza A (H1N1) Flu in Workers.

What is the best technique for washing my hands to avoid getting the flu?

Washing your hands often will help protect you from germs. Wash with soap and water or clean with alcohol-based hand cleaner*. CDC recommends that when you wash your hands – with soap and warm water – that you wash for 15 to 20 seconds. When soap and water are not available, alcohol-based disposable hand wipes or gel sanitizers may be used. You can find them in most supermarkets and drugstores. If using gel, rub your hands until the gel is dry. The gel doesn't need water to work; the alcohol in it kills the germs on your hands.

Are there medicines to treat H1N1?

Yes, however, during the current pandemic, the priority use for influenza antiviral drugs during is to treat severe influenza illness (for example hospitalized patients) and people who are sick who have a condition that places them at high risk for serious flu-related complications or are a household contact of someone at high risk for complications. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with novel H1N1 flu virus. Antiviral drugs are prescription medicines (pills, liquid or an inhaled powder) that fight against the flu by keeping flu viruses from reproducing in your body.

What does CDC recommend regarding "swine flu parties"?

"Swine flu parties" are gatherings during which people have close contact with a person who has novel H1N1 flu in order to become infected with the virus. The intent of these parties is for a person to become infected with what for many people has been a mild disease, in the hope of having natural immunity novel H1N1 flu virus that might circulate later and cause more severe disease.

CDC does not recommend "swine flu parties" as a way to protect against novel H1N1 flu in the future. While the disease seen in the current novel H1N1 flu outbreak has been mild for many people, it has been severe and even fatal for others. There is no way to predict with certainty what the outcome will be for an individual or, equally important, for others to whom the intentionally-infected person may spread the virus.

CDC recommends that people with novel H1N1 flu avoid contact with others as much as possible.

For more information on the 2009 H1N1 flu virus, call the **Lawrence-Douglas County Health Department** at (785) 843-0721, (on the web at: www.ldchealth.org), call the **Kansas Department of Health and Environment, Office of Surveillance and Epidemiology** (877) 427-7317, visit <http://www.cdc.gov/h1n1flu/> or contact your doctor, nurse or local health center.